Pancreatic cancer
Risk of diagnosis among males

SIR
- ■ Very high
- ■ High
- ■ Average
- ■ Low
- ■ Very low

New cases/year	194
Rate/100,000	11.5
Smoothed SIR distribution	
Highest	135.3
75%	103.1
Median (50%)	98.8
25%	95.9
Lowest	82.4
Geographical variation	
Evidence level	

None

(p-value 0.773)

## Level of Uncertainty

Distribution of smoothed SIR estimates according to:

### (a) Socioeconomic status

(b) Rurality

Notes: Smoothed SIR (Standardised Incidence Ratio) estimates are in comparison to the Queensland average (red line on graphs), and should not be directly compared between SLAs (Statistical Local Areas). Data are for cases diagnosed between 1998 and 2007.

Pancreatic cancer
## Risk of death within five years of diagnosis among males

5-year mortality	94.7%
Smoothed RER distribution	
Highest	126.4
75%	106.6
Median (50%)	99.2
25%	94.3
Lowest	84.1
Geographical variation	
Evidence level:	

None

(p-value 0.108)

Level of Uncertainty

### Distribution of smoothed RER estimates according to:

(a) Socioeconomic status

(b) Rurality

Notes: Smoothed RER (Relative Excess Risk) estimates are in comparison to the Queensland average (red line on graphs), and should not be directly compared between SLAs (Statistical Local Areas). Data are for 'at risk' cases in the period 1998 and 2007.

Pancreatic cancer
Risk of diagnosis among females

SIR
- ■ Very high
- ■ High
- ■ Average
- ■ Low
- ■ Very low

New cases/year	171
Rate/100,000	8.6
Smoothed SIR distribution	
Highest	120.8
75%	100.1
Median (50%)	97.4
25%	94.4
Lowest	85.0
Geographical variation	
Evidence level	

None
(p-value 1.000)

Level of Uncertainty

Distribution of smoothed SIR estimates according to:

(a) Socioeconomic status

(b) Rurality

Notes: Smoothed SIR (Standardised Incidence Ratio) estimates are in comparison to the Queensland average (red line on graphs), and should not be directly compared between SLAs (Statistical Local Areas). Data are for cases diagnosed between 1998 and 2007.

Pancreatic cancer
Risk of death within five years of diagnosis among females

RER
- ⬜ Very high
- ⬜ High
- ⬜ Average
- ⬜ Low
- ⬜ Very low

| 5-year mortality	93.7%
Smoothed RER distribution	
Highest	135.1
75%	104.3
Median (50%)	100.4
25%	96.3
Lowest	81.9
Geographical variation	
Evidence level:	

None
(p-value 0.502)

Level of Uncertainty

Distribution of smoothed RER estimates according to:

(a) Socioeconomic status

(b) Rurality

Notes: Smoothed RER (Relative Excess Risk) estimates are in comparison to the Queensland average (red line on graphs), and should not be directly compared between SLAs (Statistical Local Areas). Data are for 'at risk' cases in the period 1998 and 2007.